



Arbitrary Unitarily Invariant Random Matrix Ensembles and Supersymmetry

Thomas Guhr

III Brunel Workshop on Random Matrix Theory

Outline

- the problem and its history
- if you wish: a little bit about supersymmetry
- first step: supersymmetric representation for norm-dependent ensembles
- general case: supersymmetric representation for arbitrary rotation invariant ensembles
- some results beyond orthogonal polynomials

TG, J. Phys. A39 (2006) 12327, J. Phys. A39 (2006) 13191

The Problem and its History

Efetov's supersymmetry approach (early 80's) based on Gaussian assumption for probability densities.

physics: acceptable because of local universality

mathematics: fundamental restriction of supersymmetry ?


Hackenbroich, Weidenmüller (1995): universality proof involving supersymmetry and twofold asymptotics, not exact

Efetov, Schwiete, Takahashi (2004): superbosonization

TG (2006): algebraic duality, explicit construction

Littelmann, Sommers, Zirnbauer (2007): rigorous, threefold way

Need for Non-Gaussian Probability Densities



financial correlation matrices

empirical result deviate from Gaussian assumption

Laloux, Cizeau, Bouchaud, Potters (1999)

high-energy physics and quantum gravity, probability density:

$$P(H) \sim e^{-p - \text{tr} V(H)}, \quad V(H) = \sum_j c_j H^j$$

large-scale universality, but

/Na~~NOT~~ c~~AN~~ OBSERVING, IN ~~DO~~ OR 8g, OC

Mehta–Mahoux and Factorization

rotation–invariant probability density: $P \ H \ P \ E$

factorization: $P \ E \prod_{n=1}^N P^{(\text{ev})} \ E_n$

$$R_k \ E_1, \dots, E_k = \det \mathbf{K}_N \ E_p, E_q \quad p, q = 1, \dots, k$$

$$\mathbf{K}_N \ E_p, E_q = \sqrt{P^{(\text{ev})} \ E_p \ P^{(\text{ev})} \ E_q} \sum_{n=0}^{N-1} {}_n \ E_p \ {}_n \ E_q$$

${}_n \ E_p$ are orthogonal polF:

Supersymmetry — Linear Algebra

supervectors

$$\begin{bmatrix} z \end{bmatrix}$$

and supermatrices

$$\begin{bmatrix} a & \mu \\ b \end{bmatrix}$$

matrices a, b have commuting entries

matrices $\mu,$ have



Gaussian Integrals over Supervectors

matrix \mathbf{a} has commuting entries

$$\int e^{-\frac{1}{2} \mathbf{z}^\dagger \mathbf{a} \mathbf{z}} d\mathbf{z} = \det^{-1} \frac{\mathbf{a}}{2} \quad \text{and}$$

Supersymmetry and Gaussian Random Matrices

Gaussian ensemble of $\mathbf{N} \times \mathbf{N}$ Hermitean random matrices \mathbf{H}

k-point correlations $R_k(x_1, \dots, x_k) = \frac{1}{\prod_{p=1}^k J_p} Z_k(x, \perp J)$ $|_{J=0}$

generating function obeys the identity (yes, this is exact!)

$$Z_k(x, \perp J) = \int d\mathbf{H} e^{-\text{tr}(\mathbf{H}^2)} \prod_{p=1}^k \frac{\det(\mathbf{H} - x_p - J_p)}{\det(\mathbf{H} - x_p, \perp J_p)}$$

$$\int d\mathbf{x} e^{-\text{tr}(g^{-2} \det g^{-N} - \mathbf{x} - J)}$$

where \mathbf{x} is a $k \times k$ supermatrix

→ drastic reduction of dimensions

Posing the Problem as a Structural Issue

can we generalize this to non-Gaussian probability densities ?

is there an identity of the form

$$\int d\mathbf{H} \mathbf{P}(\mathbf{H}) \prod_{p=1}^k \frac{\det \mathbf{H} - \mathbf{x}_p - \mathbf{j}_p}{\det \mathbf{H} - \mathbf{x}_p + \mathbf{j}_p} = \int d\mathbf{Q} \det g^{-N}(\mathbf{Q}) - \mathbf{x} - \mathbf{j}$$

given an arbitrary rotation-invariant $\mathbf{P}(\mathbf{H})$, what is \mathbf{Q} ?

First Step: Norm–dependent Ensembles

consider

Examples

always $\mathbf{u} = \text{tr } \mathbf{H}^2$ and $\mathbf{w} = \text{trg } \mathbf{H}^2$

fixed trace ensemble:

\mathbf{P}^{\langle}

Arbitrary Rotation-invariant Ensembles

use bosonic fields \mathbf{z}_p and fermionic fields \mathbf{z}_p^\dagger

$$\frac{\det \mathbf{H} - \mathbf{x}_p - \mathbf{J}_p}{\det \mathbf{H} - \mathbf{x}_p + \mathbf{J}_p} = \int d\mathbf{z}_p e^{-i\mathbf{z}_p^\dagger (\mathbf{H} - \mathbf{x}_p) + \mathbf{J}_p \cdot \mathbf{z}_p}$$
$$\int d\mathbf{z}_p^\dagger e^{-i\mathbf{z}_p^\dagger (\mathbf{H} - \mathbf{x}_p) - \mathbf{J}_p \cdot \mathbf{z}_p}$$

characteristic function: $\mathbf{K} = \int d\mathbf{H} \mathbf{P}(\mathbf{H}) e^{-i\text{tr} \mathbf{HK}}$

Fourier matrix variable: $\mathbf{K} = \sum_{p=1}^k \mathbf{z}_p \mathbf{z}_p^\dagger - \sum_{p=1}^k \mathbf{z}_p^\dagger \mathbf{z}_p$

$\mathbf{P}(\mathbf{H})$ rotation invariant \longrightarrow \mathbf{K} rotation invariant

Duality between Ordinary and Superspace

introduce $N \times k$ supermatrix A , $\mathbf{z}_1 \cdots \mathbf{z}_k \quad 1 \cdots k$

$$\mathbf{K} = \sum_{p=1}^k \mathbf{z}_p \mathbf{z}_p^\dagger - \sum_{p=1}^k \mathbf{z}_p^\dagger \mathbf{z}_p \quad \mathbf{A} \mathbf{A}^\dagger$$

$$\mathbf{B} = \mathbf{A}^\dagger \mathbf{A}$$
$$\left[\begin{array}{cccccc} \mathbf{z}_1^\dagger \mathbf{z}_1 & \cdots & \mathbf{z}_1^\dagger \mathbf{z}_k & \mathbf{z}_1^\dagger & 1 & \cdots & \mathbf{z}_1^\dagger & k \\ \vdots & & \vdots & \vdots & & & \vdots & \\ \mathbf{z}_k^\dagger \mathbf{z}_1 & \cdots & \mathbf{z}_k^\dagger \mathbf{z}_k & \mathbf{z}_k^\dagger & 1 & \cdots & \mathbf{z}_k^\dagger & k \\ -\mathbf{z}_1^\dagger \mathbf{z}_1 & \cdots & -\mathbf{z}_1^\dagger \mathbf{z}_k & -\mathbf{z}_1^\dagger & 1 & \cdots & -\mathbf{z}_1^\dagger & k \\ \vdots & & \vdots & \vdots & & & \vdots & \\ -\mathbf{z}_k^\dagger \mathbf{z}_1 & \cdots & -\mathbf{z}_k^\dagger \mathbf{z}_k & -\mathbf{z}_k^\dagger & 1 & \cdots & -\mathbf{z}_k^\dagger & k \end{array} \right]$$

\mathbf{K} is $N \times N$ ordinary, but \mathbf{B} is $k \times k$ super

Equality of Invariants

for all integers $m = 0, 1, 2, \dots$ we have the identity

$$\text{tr } \mathbf{K}^m = \text{tr } \mathbf{A} \mathbf{A}^\dagger m = \text{trg } \mathbf{A}^\dagger \mathbf{A}^m = \text{trg } \mathbf{B}^m$$

non-trivial connection between ordinary and superspace

remarkable implication for characteristic function

$$\text{tr } \mathbf{K}, \text{tr } \mathbf{K}^2, \text{tr } \mathbf{K}^3, \dots \quad \text{trg } \mathbf{B}, \text{trg } \mathbf{B}^2, \text{trg } \mathbf{B}^3, \dots$$

same form as function of invariants !!

whole approach will be based on characteristic function

Spectral Decomposition

K and **B** have the same “relevant” eigenvalues !!

$$\mathbf{K} = \mathbf{V} \mathbf{\Lambda} \mathbf{V}^\dagger$$

Chain of Equalities

characteristic function satisfies

$$\mathbf{K} \quad \mathbf{Y} \quad \mathbf{y} \quad \mathbf{B}$$

alternative proof, avoiding the direct use of invariants

Construction of Generating Function

Fourier Superspace Representation

integrals over fields \mathbf{z}_p and \mathbf{p} as usual

$$\mathbf{Z}_k \propto e^{\frac{1}{2} \int d\mathbf{p} \cdot \mathbf{p} - i \text{trg}(\det g^{-N}(-\mathbf{x}^- - \mathbf{J}))}$$

arrive at a Fourier superspace representation only involving the characteristic function

$$\mathbf{Z}_k \propto e^{\frac{1}{2} \int d\mathbf{p} \cdot \mathbf{p} - i \text{trg}(\mathbf{x}^- + \mathbf{J})}$$

Generalized Ingham–Siegel–type of integral

Fourier transform of superdeterminant to power $-N$

$$\int \mathbf{d} \quad e^{-p \cdot \mathbf{i} \operatorname{tr} g} \quad \det g^{-N} - \\ \prod_{p=1}^k \operatorname{O}(\mathbf{r}_{p1}) \frac{\mathbf{i} \mathbf{r}_{p1}^{-N}}{e^{-\mathbf{p} \cdot \mathbf{r}_{p1}}} \frac{\mathbf{r}_{p2}^{N-1}}{\mathbf{r}_{p2}^{N-1}}$$

almost equal to superdeterminant to power $+N$

Probability Density in Superspace

convolution theorem in superspace yields

$$Z_k(x, \bar{x}) = \int dQ \det g^{-N} (x - Q - \bar{x})$$

desired probability density is thus Fourier backtransform

$$Q(x) = \int dP e^{i p \cdot x} \det g^{-N}$$

duality between ordinary and superspace connects Fourier transforms !!

Reduction to Eigenvalue Integrals

Fourier superspace representation has considerable advantages

\mathbf{r} and $\mathbf{I} \mathbf{r}$ invariant, apply supersymmetric Harish-Chandra–Itzykson–Zuber integral and do the group integral

$$\mathbf{R}_k(\mathbf{x}_1, \dots, \mathbf{x}_k) = \int d\mathbf{r} \mathbf{B}_k(\mathbf{r}) e^{-i\text{trg} \mathbf{x}\mathbf{r}} \delta(\mathbf{I}\mathbf{r})$$

with Berezinian (Jacobian)

$$\mathbf{B}_k(\mathbf{r}) = \det \left[\frac{1}{\mathbf{r}_{p1} - i\mathbf{r}_{q2}} \right]_{p,q=1,\dots,k}$$

The full problem is reduced to k integrals, of which k can be done trivially. This holds for arbitrary rotation-invariant probability densities $\mathbf{P}(\mathbf{H})$, including those which do not factorize!

General Result beyond Orthogonal Polynomials

another representation for correlation functions

$$R_k(x_1, \dots, x_k) = \int dH P(H) R_k^{(\text{fund})}(x - h)$$
$$h \in g(H_{11}, \dots, H_{kk}, iH_{(k+1)(k+1)}, \dots, iH_{(2k)(2k)})$$

convolution of probability density with fundT^{i120048q(d)24901071TJ /RS}

Example

probability density without factorization ($\mathbf{M}_1, \mathbf{M}_2, \dots, \mathbf{M}_k, \dots$)

$$P(\mathbf{H}) = (\text{tr } \mathbf{H}^{M_1})^{M_2} e^{-\text{tr } \mathbf{H}^2}$$

correlation functions are linear combinations of determinants

$$\mathbf{R}_k(\mathbf{x}_1, \dots, \mathbf{x}_k) = \sum_{\{m\}} \mathbf{a}_{\{m\}} \sum_{\omega} \det \left[\mathbf{C}_{m_{\omega(p)} m_{\omega(k+1)}} \mathbf{x}_p, \mathbf{x}_q \right]_{p,q=1,\dots,k}$$

$$\mathbf{C}_{m_1 m_2}(\mathbf{x}_p, \mathbf{x}_q) = e^{-\mathbf{x}_p^2} \sum_{n=0}^{N-1} \overline{\mathbf{n}}_{nm_1} \mathbf{x}_p \cdot \overline{\mathbf{n}}_{nm_2} \mathbf{x}_q$$

where

\mathbf{x}_p and $\mathbf{H} = 1000001 \text{rg} / 100 \text{cm} = \mathbf{B} \mathbf{T} \mathbf{R} \mathbf{8461000J} \mathbf{R} \mathbf{8461000J} \mathbf{B} \mathbf{T} \mathbf{R} \mathbf{8461000J} \mathbf{R} \mathbf{8461000J}$

Summary and Conclusions

- in various applications non–Gaussian probability densities
- Mehta–Mahoux theorem needs factorization
- first step: norm–dependent probability densities
- general case: arbitrary rotation–invariant probability densities
- Fourier superspace formulation only builds upon characteristic function
- all correlation functions reduced to \mathbf{k} (actually \mathbf{k}) integrals
- results beyond Mehta–Mahoux theorem
- correlation functions are convolutions involving the fundamental correlations

work in progress with M. Kieburg (Sonderforschungsbereich Transregio 12)